

120 « -10:
» -
// -
« -10: « -
120- . -2014. - .1., .1. - .179-183 »,

90-

Trichostrongylidae,

(1928)

14

8

Ostertagia, *Nematodirus* *Haemonchus*.

Nematodirus
filicolis, *Cooperia zurnabada*, *Marshallagia dentispicularis*, *Teladorsagia*
trifurcata.

Ostertagia (Ransom, 1907)

Ostertagia circumcincta (Stadgelman, 1894)

52.4%,

- 57.4

Nematodirus (Ransom, 1907)

Nematodirus oiratianus (Rajewskaja, 1929), *Nematodirus spathiger* (Railliet,
1896) *Nematodirus abnormalis* (May, 1920), *Nematodirus filicolis* (Rudolphi,

1802). *Nematodirus spathiger*. *Nematodirus*

oiratianus, *Nematodirus oiratianus* 38.1%,

36.0.

Nematodirus spathiger 21.2%, - 5.5.

Nematodirus abnormalis

Nematodirus abnormalis 5.3%,

2.2.

Nematodirus filicolis (Rudolphi, 1802)

(- -).

15.9% , - 1 , - 54.6 - 74

8.7.

41.5%,

((1960).

Trichostrongylus colubriformis (Giles, 1892, Railliet, 1898)

((1993)).

Cooperia oncophora (Railliet, 1898)

(1974).

Cooperia oncophora

3.3%, - 4.4.

4.4% , 31.6

- 1,4.

: *Cooperia zurnabada* (Antipin, 1931), *Marshallagia dentispicularis* (Assadov, 1954), *Teladorsagia trifurcata* (Ransom, 1907), *Nematodirus abnormalis* (May, 1920). *Trichostrongylus*

colubriformis (Giles, 1892), *Ostertagia ostertagi* (Stagelman, 1894; Ransom, 1907).

Trichostrongylus axei (Cobbold, 1979),

Marshallagia marshalli (Ransom, 1907), *Nematodirus helvetianus* (May, 1920),

(1979), (1974)

Cooperia zurnabada 9.5%.

- 46

127 56.8 5.4.

Marshallagia dentispicularis

4.7 %,

- 36 1.7.

Teladorsagia trifurcata (Ransom, 1907)

()

4.5% : - 54.7,

69, - 15

2.5.

Nematodirus filicolis, *Cooperia zurnabada*, *Marshallagia dentispicularis*,
Teladorsagia trifurcata

(1962), (1964), (1966),

(1961), (1961), (1959), (1976),

(1967), (1961)

()

/		%		(.)		
				.	.	.
1.	<i>Ostertagia ostertagi</i> (Stiles, 1892)	19.5	10.9	38	109	36.0
2.	<i>stertagi circumcincta</i> (Stagelman, 1894)	52.4	30.1	13	42	57.4
3.	<i>Trichostrongylus colubriformis</i> (Giles, 1892, Railliet, 1898)	3.3	4.4	82	140	133.3
4.	<i>Teladorsagia trifurcata</i> (Ransom, 1907)	4.5	2.5	15	69	54.7
5.	<i>Haemonchus contortus</i> (Rudolphi, 1803, Cobbold, 1898)	41.5	8.1	4	25	19.5
6.	<i>Marshallagia dentispicularis</i> (Assadov, 1954)	4.7	1.7	11	61	36.0
7.	<i>Marshallagia mongolica</i> (Shumacovitsch, 1938)	13.6	6.5	3	83	47.9
8.	<i>Nematodirus oiratianus</i> (Rajewskaja, 1929)	38.1	36.0	16	128	94.5
9.	<i>Nematodirus spathiger</i> (Railliet, 1896, Railliet et Henry, 1909)	21.2	5.5	15	171	25.8
10.	<i>Nematodirus abnormalis</i> (May, 1920)	5.3	2.2	16	66	41.8
11.	<i>Nematodirus filicollis</i> (Rudolphi, 1802)	15.9	8.7	1	74	54.6
12.	<i>Nematodirella longissimespiculata</i> (Romanovitsch, 1915)	3.5	1.9	15	69	54.3
13.	<i>Cooperia oncophora</i> (Railliet, 1898)	4.4	1.4	13	37	31.6
14.	<i>Cooperia zurnabada</i> (Antipin, 1931)	9.5	5.4	46	127	56.8