

« -11: = »  
« -11: ». - 2015. - . , .1. - .51-53

• ” • ”  
• •

( , , )

[1,2,3].

[4].

5 30

t 40<sup>0</sup>

6

99%

0,5-1  
30

100

20

1<sup>2</sup>

, 2 -

, 3 -

: 0-

, 4 -

, 1-

1

F<sub>1</sub>

1 -

F<sub>1</sub>

			, %		
1	79	F <sub>1</sub>	4/40	0	0
2		F <sub>1</sub>	3/30	0	0
3		F <sub>1</sub>	2/20	0	0
4	-	F <sub>1</sub>	3/40	0	0
5		F <sub>1</sub>	0	0	0
6	808	F <sub>1</sub>	0	0	0
7		F <sub>1</sub>	2/20	0	0

1

808

79

: Tr.monococcum v. vulgare, Tr.monococcum ( ), Tr.sinskayi, Tr.pseudomonococcum, Tr. urartu, Tr. araraticum, Tr.boioticum, Tr. dicoccum, Tr.Timopheevii, Tr. macha.

Tr. compactum , Tr.spelta, Tr. compactum v. rufulum.

126

60

0 (

),

Ae.ovata/Tr.durum) 32 (F<sub>2</sub> 24/ ) 6 (F<sub>7</sub>  
 0.  
 4/ 210)/Norin 10, F<sub>2</sub> (B5  
 30%.  
 F<sub>2</sub> (B5 4/ 210)/ F<sub>2</sub> (B5  
 210)/

1. . . . , 1995, . 141.
2. . . . 1968, 4, . 10-18.
3. . . . , 1983, 7, .38.

4. R. Mago, H.S. Bariana, I.S/ Dundas, G.J. Lawrence, A.J. Pryor, J.G. Ellis. Development of PCR markers for the selection of wheat stem rust resistance genes Sr24 and Sr26 in diverse wheat germplasm. Theoretical and Applied Genetics. August 2005, P. 496.