

« -11: = » -
« -11: ». - 2015. - ., .1. - .144-145

TOXOPLASMA GONDII RH

, ()
· · · , · · · , · · · ,
· · ·
,
,
·
,
[1,2].
[3,4].
(),

toxoplasma gondii RH,

,
, · · · (2005) [5].

· · · (1982) [6]

30-40

17

+4

1:51 200.

1:20
1:5 1:10

10% -
140 ,

Active T.gondi p30 protein fragment («Abcam»),
30 ,

1. Ghazy A.A., R.M. Shaapan, E. H. Abdel-Rahman. Comparative serological diagnosis of toxoplasmosis in horses using locally isolated *Toxoplasma gondii* // *Veterinary Parasitology*. - 2007. – Vol. 14, 1-2. - P. 31-36.
2. Villard O., B. Cimon, J. Franck, H. Fricker-Hidalgo, et al. Evaluation of the usefulness of six commercial agglutination assays for serologic diagnosis of toxoplasmosis // *Diagnostic Microbiology and Infectious Disease*. - 2012. - Vol. 73, 3. - P. 231-235.
3. Basso W., Hartnack S., Pardini L., Maksimov P., Koudela B., Venturini M.C., Schares G., Sidler X., Lewis F.I., Deplazes P. Assessment of diagnostic accuracy of a commercial ELISA for the detection of *Toxoplasma gondii* infection in pigs compared with IFAT, TgSAG1-ELISA and Western blot, using a Bayesian latent class approach // *Int. J. Parasitol.* - 2013. - Vol. 43, 7. - P. 565-570.
4. Al-Adhami B.H., Gajadhar A.A., A new multi-host species indirect ELISA using protein A/G conjugate for detection of anti-*Toxoplasma gondii* IgG antibodies with comparison to ELISA-IgG, agglutination assay and Western blot // *Vet. Parasitol.* - 2014. - Vol. 200, 1-2. - P. 66-73.
5. 12343.
// ;
.17.12.2002, . 12. - 4 .
6. , 1982
/ -
. (. . .) . - 15 .