

« -11: = » -  
« -11: ». - 2015. - . , .1. - .208-210

-  
.. „ .

.  
,  
44-50% 27% 30%

( , , , , )  
- , -  
10-15%

-  
, , - ,  
[1].

( )  
Casas E. 14 [2].

[3].  
36-40 4-6  
13-16

[4].  
« - »

2014

6  
3

5

24

15,86 × 14,5% (P >0,99),  
 2,41% 5,27%, (P >0,95).  
 10,18% (P >0,99). 13,91%,  
 10,59 × 8,95%

[5]

A. Ali

(P >0,95).

(P >0,95).  
(P >0,95),

×

×

-

«

»

-

1

85-

, 2014. 78-81 .

2 Casas E., Freking B.A., Leymaster A. Evaluation of Dorset, Finnsheep, Romanov, Texel, and Montadale breeds of sheep: V. Reproduction of F1 ewes in spring mating seasons. *Journal of Animal Science*, 2005. 83: 2743-2751 p.

3 Scales G.H., Bray A.R., Baird D.B., O'Connell, T.L. Knight. Effect of sire breed on growth, carcass, and wool characteristics of lambs born to Merino ewes in New Zealand. *New Zealand Journal of Agricultural Research*, 2000. 43: 93-100.

4

2- , . - ., 2011. 190-201 .

5 Ali A., Morrical D.G., Hoffman P., Berger P.J. Pas. Evaluating Texel-, Suffolk-, and Columbia-Sired Offspring: II. Post weaning Growth and Carcass Traits Under Feedlot and Pasture Feedlot Finishing Systems<sup>1</sup>. *The Professional Animal Scientist*, 2005. 21:434-442

: . . . , . . . . ,