

·
2013. - 1 (76). - .42-52

/

· , . .



· , : ,
() .

· ,

· , .

· : , , ,

· ,

· , [1-7].
· ,
· ,

[8-10].

: 150

10%

[11-12],

[1,13-15].

，
-

(
)，
，
(
)
)，
(
)，
(
).

：
()。
：

() - 300 500 -

，
，
，
-
，
，
：
-

100
XX

9

(1909-1912; 1924-1927; 1931-1933;
1944-1947; 1953-1956; 1967-1970;
- 1977-1982; 1988-1991; 1997-2003 .)
[2,5,6,7,16,17].

(10-12)

()

().

(70-75%)

(
49,5° – 52° . . . 61,5° – 80° . . .)
[1,8,10].

« »

« » [18].

(, , , , ,) , « » « »

, - . , () , , . - . « » « » 1960- .

- -

1950- - 1960- . , , , , , 25 . (1976 -

1986 .) 11-12 . (1997 – ,
 1999),
 - , ,
 . -
 , , . ,
 , ,
 , , ,
 . - .
 ,
 29 / 1986 0,1 / -
 2000 , , 1997
 . 2003 ,
 ,

1997-2003 .

50
1997-1998

1000-1200
[2,5,7].

3000 – 5000

..

(140

[2,5,7].

11-12 18-19

2,5-4,5 /

6-8 / ,

- 28-36

3

,7-8

, 10-11

,12-13

40-60

,
 .
 2
 ,
 -
 -
 [2,4,5].
 ,
 ,
 ,
 1999-2000 . 4 ,
 : - .
 (60 .), (-
 30 .), - ,
 (18 .), - ,
 (5 .). , -
 .
 ,
 .
 1 2 ,
 -
 5200 180000-200000 ,
 1 2 -
 .
 .
 :
 ,
 . ,
 . ,
 .

1, 1999 () 1997-1998 2, 2,5 (1\$=131) 220

.....

2000 2,8
(20,1
)

.....

400

"

2000 - 8,1
2001 - 4,8
5
(

.....

: 2002 - 1,2
2003 - 601,5 , 2004 - 506
[19].

"
«

»

.....

()

80-120
2000-2001 (100-300).
13
3,5

95-99%.

[1,10].

12,9 2000-2001 10 5
5 20-40 2012

2004 506 [20].
2005 2011
2228,4 7
4,4

2012

2010-2011-2012 .

.),

(

).

2012

1883,4

258,3

«

»

2012

:

10-14

20

100

1²,

2000-4000

1²:

7-10

;

(

,

-

1997-2003 .

[10].

[1,8,10,19,21].

GPS/

(),

(,),

(,).

- 2001

, «

»

15-20

[10].

14. // -
- 10- , 1993: . - . - .87-88.
15. / . . ,
16. . . (.) . - : « » , 2003.- 48 .
- .25-38. // . - , 2001. -
17. // . - .1. - , 2002. - .58-66.
18. : , 1970. -271 .
19.
- // . The phytosanitary safety of agroecosystems. - , 2010. - . 5-13.
20. (Calliptamus italicus L.) 2012 // . 55
- " : , , " , 2012. - .241-243.
21. « ()» . - , , 12-16 2012.

Summary

Particularly hazardous is Asian, Italian and Moroccan locusts. Areas of mass reproduction are located in many regions of the country. While some of the most effective among them recognized chemical. The analysis of the existing methods for massive chemical treatments over large areas in the midst of outbreak of locusts destabilizing the ecological situation have been carried out. As an alternative a proactive approach, ensuring long-term sustainable solution to the locust problem was proposed. This strategy is the result of many years of research, which has a solid scientific basis, and confirmed by extensive practice.